

A neural network approach to estimate water-column nutrient concentrations and carbonate system variables in the Mediterranean Sea: **CANYON-MED**

Marine Fourrier¹, Laurent Coppola¹, Fabrizio D’Ortenzio¹, Hervé Claustre¹

(1) Sorbonne Université, CNRS, Laboratoire d’Océanographie de Villefranche, LOV, F-06230 Villefranche-sur-Mer, France.

Contact:
fourrier@obs-vlfr.fr




1– INTRODUCTION

The Mediterranean Sea (MedSea) is characterized by a short residence time¹, an elevated **salinity** and is one of the largest oceanic nutrient-depleted areas² exhibiting an eastward-increasing **oligotrophy** gradient³.


In the context of a critically undersampled ocean, the development and intensive use of instrumented *in situ* **autonomous platforms** allows to densify the measurements of some biogeochemical variables (remaining however far from exhaustive).

2– CANYON-MED METHOD

We compiled a **homogeneous quality-controlled** dataset of *in situ* measurements from 1981 to 2018, including samples of core variables such as dissolved oxygen (O₂), temperature (T), salinity (S) as well as **macronutrients** (nitrates: NO₃⁻, phosphates: PO₄³⁻, silicates: Si(OH)₄) and **carbonate system variables** (total alkalinity: A_T, total carbon: C_T, and *in situ* pH on the total scale: pH_T).



Map of the 35 cruises and the time-series used for the training and validation of the CANYON-MED neural networks. (~8 000 values for NO₃⁻, ~11 000 for PO₄³⁻ and Si(OH)₄; and ~6 000 for A_T and pH_T, ~4 000 values for C_T).



Schematic representation of the CANYON-MED neural networks. Day: day of year
Adapted from ^{4, 5}

The **CANYON-MED** (for Carbonate system and Nutrients concentration from hYdrological properties and Oxygen using a Neural-network in the MEDiterranean Sea; in prep) **neural network-based** method provides estimations of **nutrients** and **carbonate system** variables from systematically measured oceanographic variables (T, S, O₂, geolocation, sampling date), such as the ones measured by profiling floats.

Statistics (coefficient of determination (R²), Mean Absolute Error (MAE) and Root Mean Squared Error (RMSE)) of the regressions between the CANYON-MED retrieved values with the corresponding *in situ* measurements.

	CANYON-MED		
	R ²	MAE	RMSE
NO ₃ ⁻ (μmol.kg ⁻¹)	0.95	0.47	0.77
PO ₄ ³⁻ (μmol.kg ⁻¹)	0.91	0.028	0.047
Si(OH) ₄ (μmol.kg ⁻¹)	0.96	0.40	0.70
A _T (μmol.kg ⁻¹)	0.95	7	11
C _T (μmol.kg ⁻¹)	0.89	9	14
pH _T	0.86	0.010	0.016


In order to better understand nutrients and pH dynamics, we test CANYON-MED’s predictions on **3 BGC-Argo floats** equipped with **nitrates** and **pH** sensors, in contrasted areas of the **MedSea** (North-Western Mediterranean Sea, Adriatic and Ionian Seas, Levantine Basin), along the W-E oligotrophy gradient.

3– APPLICATION ON MEDSEA BGC-ARGO FLOATS

For each panel: Timeseries of NO₃⁻ float data, CANYON-MED NO₃⁻, CANYON-MED PO₄³⁻, CANYON-MED Si(OH)₄, pH_T float data, CANYON-MED pH_T, CANYON-MED A_T and CANYON-MED C_T. Location of the BGC-Argo float, the red star and yellow square representing the location of the 1st and last profiles.

NW MEDITERRANEAN


6902969



Good agreement between float and CANYON-MED NO₃⁻.
For pH_T, small difference in surface values. PO₄³⁻, Si(OH)₄, A_T and C_T coherent with bottle data.
→ Dynamics well reproduced.

ADRIATIC & IONIAN


6901865



Good agreement between float and CANYON-MED NO₃⁻.
pH_T, PO₄³⁻, Si(OH)₄, A_T and C_T coherent with bottle data.
→ Satisfying results in relatively coastal waters!

LEVANTINE BASIN

6902904



Similar tendencies between float and CANYON-MED NO₃⁻.
pH_T, PO₄³⁻, Si(OH)₄, A_T and C_T in the same range as bottle data.
→ Promising results in under-represented area of CANYON-MED’s training.

4 – CONCLUSIONS AND PERSPECTIVES

The regional CANYON-MED neural networks produce satisfactory results: accuracies of **0.77**, **0.047**, and **0.70 μmol.kg⁻¹** for NO₃⁻, PO₄³⁻ and Si(OH)₄ respectively, and of **0.016**, **11 μmol.kg⁻¹** and **14 μmol.kg⁻¹** for pH_T, A_T and C_T respectively, comparable to the ones obtained by the NO₃⁻ (±1 μmol.kg⁻¹) and pH_T (±0.005) sensors mounted on BGC-Argo floats. Applied to the large Argo network, CANYON-MED will contribute to **Quality Control** and **Delayed Mode** processing for NO₃⁻ and pH_T as well as to increase the amount of biogeochemical data in the MedSea, dramatically improving our understanding of nutrients, pH and pCO₂ variability. Furthermore, CANYON-MED will generate “virtual” data of parameters not yet measured by BGC-Argo floats (namely PO₄³⁻, Si(OH)₄, A_T, and C_T).

References: /1/ Somot, S., Sevault, F., & Déqué, M. (2006). Transient climate change scenario simulation of the Mediterranean Sea for the twenty-first century using a high-resolution ocean circulation model. *Climate Dynamics*, 27(7–8), 851–879. DOI: 10.1007/s00382-006-0167-z /2/ Ignatiades, L. (2005). Scaling the trophic status of the Aegean Sea, eastern Mediterranean. *Journal of Sea Research*, 54(1), 51–57. DOI: 10.1016/j.seares.2005.02.010 /3/ Pujo-Pay, M., Conan, P., Oriol, L., Cornet-Barthaux, V., Falco, C., Ghiglione, J.-F., et al. (2011). Integrated survey of elemental stoichiometry (C, N, P) from the western to eastern Mediterranean Sea. *Biogeochemistry*, 8(4), 883–899. DOI: 10.5194/bg-8-883-2011 /4/ Sauzède, R., Bittig, H. C., Claustre, H., Pasqueron de Fommervault, O., Gattuso, J.-P., Legendre, L., et al. (2017). Estimates of Water-Column Nutrient Concentrations and Carbonate System Parameters in the Global Ocean: A Novel Approach Based on Neural Networks. *Frontiers in Marine Science* 4. DOI: 10.3389/fmars.2017.00128 /5/ Bittig, H. C., Steinhoff, T., Claustre, H., Fiedler, B., Williams, N. L., Sauzède, R., et al. (2018). An Alternative to Static Climatologies: Robust Estimation of Open Ocean CO₂ Variables and Nutrient Concentrations From T, S, and O₂ Data Using Bayesian Neural Networks. *Frontiers in Marine Science* 5. DOI: 10.3389/fmars.2018.00328.

Acknowledgments:

This work was supported by the E-ARISE H2020, WG4.

